

Research

Embodied cognitive evolution and the cerebellum

Robert A. Barton*

Evolutionary Anthropology Research Group, Department of Anthropology, Durham University, Dawson Building, South Road, Durham DH1 3LE, UK

Much attention has focused on the dramatic expansion of the forebrain, particularly the neocortex, as the neural substrate of cognitive evolution. However, though relatively small, the cerebellum contains about four times more neurons than the neocortex. I show that commonly used comparative measures such as neocortex ratio underestimate the contribution of the cerebellum to brain evolution. Once differences in the scaling of connectivity in neocortex and cerebellum are accounted for, a marked and general pattern of correlated evolution of the two structures is apparent. One deviation from this general pattern is a relative expansion of the cerebellum in apes and other extractive foragers. The confluence of these comparative patterns, studies of ape foraging skills and social learning, and recent evidence on the cognitive neuroscience of the cerebellum, suggest an important role for the cerebellum in the evolution of the capacity for planning, execution and understanding of complex behavioural sequences—including tool use and language. There is no clear separation between sensory–motor and cognitive specializations underpinning such skills, undermining the notion of executive control as a distinct process. Instead, I argue that cognitive evolution is most effectively understood as the elaboration of specialized systems for embodied adaptive control.

Keywords: brain; neocortex; cerebellum; evolution; cognition; language

1. INTRODUCTION

The idea that there was likely to have been a wide variety of selection pressures on cognitive abilities, and a corresponding variety of neural evolutionary responses [1–3], has been rather lost in the current enthusiasm for monolithic explanations for the evolution of large brains, including social intelligence [4], behavioural flexibility [5] and general intelligence [6,7]. These general explanations are associated with the search for a single comparative brain measure that best reflects cognitive ability, such as neocortex ratio [8,9], ‘executive brain’ ratio [10,11] and even whole brain size [12,13]. A relatively strong correlation between the putatively critical behavioural variable and a particular comparative brain measure is sometimes taken to suggest that the measure identified does indeed most effectively capture the neurological basis of cognitive evolution [8,13].

Empirically, there is a problem with this approach: comparative studies have not produced a single, unified picture of the relationship between such measures and behaviours. Healy & Rowe [14, p. 456] summarized the picture as one of a ‘bewildering array of correlations between brain size and behavioural

traits’, a picture which shows little sign of resolving. For example, while Dunbar & Shultz [9] argue that the central aspect of primate brain evolution is the correlation between neocortex size and social group size, Reader *et al.* [11] find that neocortex and ‘executive brain’ size correlate strongly with a composite measure of general intelligence that cuts across the social/non-social domain, and that this composite measure does not correlate with social group size.

There are also theoretical reasons to question the underlying assumption that intelligence evolved in a unitary way and can in principle be measured by a single, ideal comparative brain measure. First, which measure achieves the strongest correlation with a putatively important aspect of behaviour should not be the *sine qua non* for deciding how to measure cognitive evolution. Indeed, it is circular to argue that a particular measure is ideal because it most strongly supports a hypothesis. Second, organisms are subject to a wide variety of challenges. For example, they may be aquatic or terrestrial; they may be active at night or by day; they may be more or less social; they may graze on abundant plants, search for rare fruits, or hunt for prey; they may learn complex songs; they may store food and recover it by memory. Each of these and other dimensions of behavioural ecology has been shown to correlate with the brain size and/or with a specific and relevant aspect of brain structure [14–20]. And studies of phylogenetic variation in the brain structure of mammals and birds indicate not one or two dimensions of variation but many [21–24].

*r.a.barton@durham.ac.uk

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rstb.2012.0112> or via <http://rstb.royalsocietypublishing.org>.

One contribution of 15 to a Theme Issue ‘New thinking: the evolution of human cognition’.

A further problem is that critical assumptions underlying the use of brain size indices remain largely untested. The volume of a brain region is potentially related to cognitive capacities to the extent that it correlates with more functionally meaningful variables such as numbers of neurons and synapses. Recent works suggest that the relationship between volume and neuron number or density varies between taxonomic groups and between brain structures [25,26]. Such variability potentially presents problems for inferring functional consequences from relative size measures such as volumetric ratios between one structure and another. Here I examine the consequences of volumetric ratios for relative numbers of neurons in the neocortex and cerebellum, and I argue that an excessive emphasis on the neocortex has obscured important patterns in brain evolution and led to an unwarranted neglect of the cerebellum. I then re-examine phylogenetic correlates of neocortex and cerebellum size.

In the light of these results, I develop a synthesis of the comparative, anatomical and functional neuroscience data. This synthesis stresses the unity of sensory–motor and cognitive evolution. Classically, distinctions are made between cognition, as a process of interpreting and integrating information about the outside world, the perceptual information that this process is about, and the motor commands that represent the output of cognitive processes [27]. More recently, these distinctions have been broken down by the recognition that cognition is best conceived as a set of processes mediating the adaptive control of bodies in environments: the concept of embodied cognition [28–33]. This perspective suggests that ‘a key aspect of human cognition is... the adaptation of sensory-motor brain mechanisms to serve new roles in reason and language, while retaining their original function as well.’ [34, p. 456]. Here I argue that understanding brain evolution both contributes to and is benefited by this perspective.

2. METHODS

I use phylogenetic comparative analyses of brain component volumes and neuron numbers to test hypotheses about the evolutionary determinants and cognitive consequences of brain structure evolution. Analyses include broad patterns of brain evolution across mammalian orders and more focused analyses of behavioural correlates within primates. In the absence of direct observation of evolutionary processes, phylogenetic comparative analysis provides a powerful technique for investigating evolutionary patterns and processes [35] such as correlated trait evolution. A variety of methods now exist, but the underlying rationale of each is that combining information on phylogenetic relationships among species with data on their phenotypic traits allows one to statistically model the evolution of those traits along the branches of the tree representing their relationships [35]. To assess how different brain and behavioural traits evolved in relation to one another, I used phylogenetic generalized least squares, which incorporates phylogeny into statistical models [36–38]. Further details of this method and data used are provided in the electronic supplementary

material. Results are presented in the context of discussion of a series of key questions, and embedded where appropriate to the discussion rather than consolidated in a single results section.

3. IS THE NEOCORTEX THE ‘INTELLIGENT’ BIT OF THE BRAIN?

The brain structure most often identified with ‘higher’ cognitive functions is the neocortex [39], having been described, for example, as ‘the crowning achievement of evolution and the biological substrate of human mental prowess’ [40]. The assumption that the neocortex is the place to look for evidence about cognitive evolution drives much comparative research and even the selection of regions of interest in the study of fossil hominin endocasts [41].

Why this focus on the neocortex? One reason is undoubtedly the simple observation that it is disproportionately large in large-brained species. In small-brained mammals such as shrews the neocortex comprises as little as 15 per cent of brain volume, whereas in monkeys the corresponding figure is about 65–75 per cent and in humans it is about 80 per cent [42,43]. The correlation between brain size and neocortical proportion (or ratio) may, however, have more to do with allometric scaling than with cognitive selection pressures. Cortical proportions are generally high in large-bodied species such as sea lions (66%) [44], camels (71%) [45] and sperm whales (87%) [45]. Whilst it might be tempting to speculate on the hitherto unappreciated intelligence of these species, the most parsimonious explanation is that they are just large animals. Indeed, controlling for phylogenetic effects, there is a strong correlation between body size and proportion of the brain that is neocortex (phylogenetic least squares (PGLS); $\lambda = 0.92$, $t = 14.23$, $p < 0.0001$). There is no such correlation for the cerebellum ($\lambda = 0.93$, $t = 1.25$, $p = 0.21$).

Why does the cortex balloon in proportional size as body size (and overall brain size) increase? Apparently because of a need to devote increasing brain space to making cortical connections: larger cortices are increasingly made up of white rather than grey matter (figure 1*a*, see also [46,47]). In the cerebellum, there is a much less steep increase in white matter volume with overall size (figure 1*b*; and see [47]). Hence connectivity scales in different ways in these two structures.

The reasons for this difference in white versus grey matter scaling presumably relate to the basic connective architecture of the mammalian brain. Much of the neocortical white matter consists of fibres that make long-range connections, in which increases in axon diameter and myelination are necessary to preserve processing speed over longer conduction distances in larger brains [48,49]. The relative ballooning of the neocortex in large (and large-brained) animals may therefore be driven by allometric connective constraints rather than by any special cognitive selection pressures. One implication is that the ratio measures of relative brain structure size used commonly in comparative studies, such as neocortex ratio [8], ‘executive’ brain ratio [7,10,11] and ‘cerebrotype’ [50] conflate allometric scaling with selection on specific




Figure 1. White matter proportion increases more steeply with size in neocortex than in cerebellum. The proportion of volume of (a) neocortex and (b) cerebellum that is white matter, plotted against volume of each structure (mm^3). The graphs plot data for the same species and the PGLS slopes are significantly different (see text).




Figure 2. Contrast in the pattern of variation in the proportion of the brain composed of neocortex versus cerebellum when expressed as (a) volume proportion and (b) proportional number of neurons. Dark bars represent cortical proportions and light bars denote cerebellar proportions.

brain regions. A volumetric ratio between neocortex and other structures potentially underestimates selection on non-cortical (e.g. cerebellar) functions.

The striking variation in the proportional size of the mammalian neocortex cannot therefore be simplistically read as reflecting selection specifically on cortical functions. This is further emphasized by the lack of correspondence between volumetric ratios and numbers of neurons. In stark contrast to the way that cortical volume proportion scales up with brain size, cortical neuron number proportion is unrelated to brain size [26] and unrelated to cortical volume proportion [25]. Similarly, the ratio of cortical to cerebellar volumes is uncorrelated with the ratio of cortical to cerebellar neurons (PGLS; $\lambda = 0.63$, $t_{2,23} = 1.13$, $p = 0.27$), casting doubt on the functional significance of volumetric ratios. Neuron density decreases as brain size increases in both neocortex (PGLS: $\lambda = 0.83$, slope = -0.23 , $t_{2,23} = 4.55$, $p < 0.0002$) and cerebellum (PGLS: $\lambda = 0.76$, slope = -0.04 , $t = 2.43$, $p = 0.02$), but the decline is significantly steeper in the neocortex (difference in PGLS coefficients: $t = 3.92$, $p = 0.0008$). The same is true when neuron densities of the two structures are related to their volumes rather than to overall brain size ($t = 2.86$, $p = 0.009$). Hence, the increase in neocortical volume proportion with brain size is traded off against a steeper decrease in neuron density.

Evidently there are different scaling constraints on each structure. Figure 2 illustrates the markedly

different patterns of cross-species variability in proportional volumes and proportional neuron numbers, as well as the much larger number of neurons in the cerebellum than in the neocortex of all species. These results question both the validity of volumetric ratios as useful measures of information-processing capacity and the justification based on their variability across species for the near-exclusive focus of comparative studies on the neocortex.

As pervasive as the assumption that neocortical expansion underpinned the evolution of 'higher' cognition is the assumption that it was the frontal lobes in particular that expanded most. Comparative data are relatively sparse, and most attention has focused on whether human frontal lobes are relatively large compared with their size in other primates [51–60]. The question has until recently remained unresolved, largely because of confusion over whether the proportional size or the size relative to allometric scaling provides the most useful measure. Because frontal lobe volume, like overall neocortex volume but to an even greater extent, scales hyper-allometrically, human frontal areas are large as a proportion of brain or neocortex size [53,54,59,60]. However, there is no more reason to think that proportional or absolute volume is a good measure of functional specialization for the frontal lobes than there is to believe it for the neocortex as a whole. Recent allometric analyses reveal that, although absolute and proportional frontal region size increased




Figure 3. Difference in relative numbers of neurons in (a) the neocortex and (b) cerebellum of primates (open circles) compared to other mammals (filled circles). Controlling for numbers of neurons in the rest of the brain, the difference between primates and non-primates is significant for neocortex (PGLS; $\lambda = 0.86$, $t_{3,23} = 3.43$, $p = 0.002$) and cerebellum (PGLS; $\lambda = 0.76$, $t_{3,23} = 4.54$, $p = 0.0002$). The effect is stronger for cerebellar neurons and the primate–non-primate difference in cerebellar neurons is still near-significant after controlling for neocortical neurons (PGLS; $\lambda = 0.61$, $t_{4,23} = 2.02$, $p = 0.06$).

rapidly in hominins, this change was associated with size increase in other areas and whole brain size, rather than with specialization for enlarged frontal lobes specifically [57,61–63]. Consistent with allometric effects, neuron densities are particularly low in human frontal cortex [58]. Interestingly, there is stronger evidence for relative enlargement of temporal lobe structures [64,65]. This does not suggest that the frontal lobes were unimportant in cognitive evolution, just that their importance needs to be interpreted in terms of the areas with which they connect and with which they have co-evolved, including the cerebellum [61,62].

4. CEREBELLA COMES TO THE BALL: RELATIVE EXPANSION AND CO-VARIATION OF NEOCORTEX AND CEREBELLUM

Although allometric scaling explains much of the variation in proportional neocortex size, it does not explain all of it. After taking scaling against other brain structures into account, primates have relatively large neocortices [23], and a relatively high density of cortical neurons [48]. However, the cerebellum is also larger [66] and contains more neurons in primates compared to other mammals (figure 3). This conjoint expansion of the two structures early in primate




Figure 4. Correlated evolution of neocortex and cerebellum size in mammals. Neocortex size and cerebellum size are positively correlated after controlling for phylogenetic effects and volume of other brain regions (PGLS, neocortex volume regressed on volume of cerebellum controlling for volume of the rest of the brain; $\lambda = 0.97$, $t_{3,298} = 8.85$, $p < 0.0001$).

evolution reflects a general evolutionary trend for the two structures to evolve together, in primates in particular [23,26,62,67], and more generally during mammalian evolution (figure 4).

There are three compelling aspects of the evidence for correlated evolution of the neocortex and cerebellum. First, it is apparent after accounting for variability in the size of other brain structures, discounting the possibility that it is a reflection of some global allometric or developmental constraint acting across the whole brain. Second, there is a striking correspondence between the patterns of correlated evolution among specific components of the cortico-cerebellar system and their anatomical connectivity, down to the level of individual nuclei [62,67]. Third, it is evident not just in terms of volumes, but also in two independent data sets on numbers and densities of neurons (figure S1 in the electronic supplementary material).

The linkage between neocortical and cerebellar expansion suggests that both contributed significantly to brain size evolution. Indeed, a phylogenetic analysis reveals that, controlling for body mass, mammalian brain size is positively and independently correlated with both neocortex and cerebellum, and also that there is a significant interaction between the effects of the two structures on brain size (PGLS, brain mass regressed on: body mass, $t = 8.47$, $p < 0.0001$; neocortex, $t = 19.73$, $p < 0.0001$; cerebellum, $t = 12.35$, $p < 0.0001$; interaction between neocortex and cerebellum, $t = 4.04$, $p < 0.0001$; $\lambda = 0.92$, $n = 298$ mammal species). The combination of significant main and interaction effects suggests that the evolution of brain size was a product of both independent and co-ordinated size change of neocortex and cerebellum.

Previous work demonstrated a strong association between relative neocortex size and visual specialization in non-human primates [19,20,48]. Is the pattern of cortico-visual evolution confounded by cortico-cerebellar evolution? Further analysis suggests not: neocortex volume is significantly and independently correlated with volumes of both visual thalamus (LGN) and cerebellum, after accounting for variation in other brain structures (PGLS,

neocortex volume regressed on volumes of cerebellum, LGN and rest of the brain; $\lambda = 0.87$, $r^2 = 0.98$; LGN, $t_{4,42} = 3.46$, $p = 0.001$; cerebellum, $t_{4,42} = 4.20$, $p = 0.0002$). The same pattern is found after subtracting primary visual area V1 from total neocortex volume ($\lambda = 0.89$, $r^2 = 0.98$, $n = 42$; LGN, $t_{4,42} = 2.82$, $p = 0.008$; cerebellum, $t_{4,42} = 4.26$, $p = 0.0001$), emphasizing that extra-striate cortex is not ‘non-visual’ [68]. The latter point is important, as different scaling trends for V1 and non-V1 against brain size have been misinterpreted as evidence against the visual specialization hypothesis [59]. In summary, variation in primate neocortex size is strongly related to the evolution both of visual structures and the cerebellum.

Several comparative studies suggest that cerebellar expansion, specifically involving the lateral cerebellum, was especially marked in apes [69–71]. It therefore seems that the cerebellum—modestly concealed beneath the volumetrically dominating neocortex, and largely ignored—may be the Cinderella of the study of brain evolution. This conclusion is reinforced by growing evidence that ascribing to it the task of basic chores in adaptive neural processes has also been a mistake.

5. COGNITIVE IMPLICATIONS

It has long been known that the cerebellum is involved in sensory–motor control and learning of motor skills [72,73]. The relative expansion of the cerebellum in primates together with stereopsis and elaboration of the visual system [19,20,68] presumably underpins primates’ fine visuo-motor control and manual dexterity. For example, smooth-pursuit eye-movements in primates are based on a unique cortico-cerebellar pathway that evolved together with foveal vision [74].

However, in the past 10 years or so considerable evidence has accumulated that the cerebellum has a broader role than previously recognized, including emotion [75,76], non-motor associative learning [77], working memory and mental rehearsal [77,78], verbal working memory and other language functions [76,78–81], spatial and episodic memory [79,81,82], event prediction [83], empathy and predicting others’ actions [84–87], imitation [88], planning and decision-making [79,89,90], individual variation in cognitive performance [91], and cognitive developmental disorders including autism [80,92].

Some have argued that the case for cognitive functions of the cerebellum remains unproven [72,93]. The details of this debate are beyond the scope of this paper, but three general points can be made. First, although some studies have been criticized for failure to control for eye movements [93], the overall weight of evidence of many clinical and functional imaging studies indicates cerebellar involvement in a wide variety of cognitive processes [94]. Second, the cerebellum and neocortex are massively interconnected [78,90], and these connections involve many cortical areas, again suggesting a wide range of functions. Third, the distinction between sensory–motor control and cognition is arbitrary and an impediment to understanding brain function and evolution. Dissolving this distinction makes the debate on the cerebellum one about the

range of its functions rather than a question of whether or not it has cognitive functions.

The classical view of cortico-cerebellar connections was that the cerebellum collected sensory information and returned it to primary motor cortex for the generation of movements [90]. However, it is now known that all major cortical regions, i.e. beyond motor cortex and including frontal and prefrontal areas, have reciprocal connections with the cerebellum. These cortico-cerebellar loops form multiple, independent anatomical modules which are architecturally quite uniform [90,95]. This anatomical uniformity together with functional data suggests basic similarities in the computations performed in different functional domains by different cortico-cerebellar modules [95,96]. These computations act as internal models or simulations of cortical processes that continuously update and error-correct responses, based on a comparison of actual and expected inputs, and they underlie a wide range of behavioural control processes [89,95,96]. Thus, internal models generated by the cerebellum guide behaviour in different domains. Direct control of behaviour, prediction of its consequences and reasoning about it may be mediated by similar cortico-cerebellar computations, with functional differences determined by which specific cortico-cerebellar modules are activated and their connectivity with other systems. Simulations computed ‘offline’ (as in the planning of sequences of behaviour), and those generated by observing other individuals (allowing prediction of their behaviour), are widely considered to be ‘cognitive’, or ‘executive’ processes. However, essentially the same kinds of computation appear to underlie sensory–motor and more ‘cognitive’ control processes [95,96], including speech [97].

6. ADAPTIVE NEURAL CONTROL PROCESSES CUT ACROSS DOMAINS, USE SIMILAR COMPUTATIONS AND SHARE CIRCUITS

Computational commonality across functional domains with overlapping neural substrates may in fact be a rather generic feature of the brain. For example, social and non-social decision-making activate adjacent brain regions in the anterior cingulate and are mediated by the same computational processes, suggesting that social and non-social cognition may not be as encapsulated or specialized as has been assumed [98]. In another example, social rejection and physical pain activate overlapping brain regions, including somatosensory cortex and cerebellum [99]. Similarly, Shackman *et al.* [100] argue that cognitive control, negative affect and pain share an overlapping neural substrate and a common computational structure, and suggest the term ‘adaptive control’ as an encompassing term for these processes. Shackman *et al.* [100] point to the intriguing fact that all three processes activate muscles of the upper face, further emphasizing commonalities across processes traditionally distinguished as ‘executive’ and ‘non-executive’. Here, functional distinctions result from divergent patterns of connection rather than fundamentally different types of computation. Thus, individual brain regions contribute to multiple functional

Table 1. Phylogenetic generalized least squares analysis of the relationship between volumes of brain components and behavioural variables. Significant associations indicated in bold. In model 1, whole brain size was regressed on body mass, group size and extractive foraging. In models 2 and 3, volumes of the individual brain regions were treated in the same way as in model 1, but the volume of the residual portion of the brain (brain – (neocortex + cerebellum)) was included as a predictor variable. Hence, these results indicate significant relationships between behavioural variables and size variation of neocortex and cerebellum relative to the size of the rest of the brain.

model parameter	model 1. whole brain size $t_{4,42}$, p -value	model 2. neocortex $t_{4,42}$, p -value	model 3. cerebellum $t_{4,42}$, p -value
body mass	18.0, <0.0001	0.95, 0.35	3.12, 0.003
volume of residual brain portion	—	12.37, <0.0001	8.93, <0.0001
group size	3.47, 0.001	5.55, <0.0001	2.64, 0.012
extractive foraging	2.73, 0.01	2.07, 0.045	3.58, 0.0009
λ	>0.99	>0.99	>0.99
model summary			
maximized log-likelihood	38.7	33.6	65.2
adjusted R^2	0.92	0.98	0.99

modules, and become secondarily adapted for use in different systems through the evolution of new connections [32,101].

7. TECHNICAL SKILLS, COGNITIVE SEQUENCING AND LANGUAGE

An adaptive control function in which the cerebellum plays a critical role is the modelling, prediction and organization of sequences of events and behaviours, including sequences involved in tool-making and use, and language comprehension and production [73,77, 78,81,90,97,102]. Thus, the cerebellum is involved in learning of procedural sequences, recognition of correct spatial and temporal relations among behaviourally relevant actions, temporal organization of verbal utterances and planning of speech, and mental rehearsal [81]. It also seems to be involved in processing more abstract sequences such as in story comprehension [103].

There is an intriguing confluence between this evidence for cerebellar involvement in the temporal organization, comprehension and learning of sequences, evidence of cerebellar expansion in great apes [69–71], and observations of the facility of these species for extractive foraging and tool use [104], including the flexible recombination of tool components or elements of a problem [105], and for solving problems requiring sensitivity to sequence information [106]. Byrne [107,108] argues that great ape extractive foraging skills are based on iterated, hierarchically organized, multi-stage algorithms for solving ‘syntactical’ problems (problems requiring behaviours to be performed and flexibly recombined in functional sequences), and that they are socially learned, possibly by programme-level imitation [109]. Cerebellar specialization in ancestral great apes may therefore have been a precursor to neural capacities underlying the later development of cumulative cultures of more complex technologies in hominins [110,111].

Parallels between the organization of behavioural sequences in extractive foraging and tool use, on the one hand, and in language processing, on the other hand, may indicate that neural specialization for the first was a pre-adaptation for the second [101,112–114], with

gestural communication probably representing an intermediate stage [114]. Indeed, there is overlap in brain areas activated during linguistic processing and other hierarchically organized motor acts such as tool construction [32,101,112,113]. In addition to classical cortical language areas, the cerebellum is activated by speech comprehension tasks [97,101,115]. Hence, language may have been built from pre-existing sensory–motor specializations common to all great apes [101].

8. TECHNICAL VERSUS SOCIAL INTELLIGENCE AND BRAIN EVOLUTION

The evidence of cerebellar expansion and involvement in diverse cognitive functions suggests that the well-known link between neocortex size and social group size [8] may not be the only important feature of primate neuro-cognitive evolution; selection on foraging skills may have been important too [70,116]. A new phylogenetic comparative analysis controlling for allometric effects supports this contention (table 1). First, the well-known correlation between neocortex (or brain) size and social group size is recovered, but neocortex size also correlates with foraging skills. Second, cerebellum size also correlates with both types of behavioural variable. Third, there is evidence of an evolutionary brain–behaviour double dissociation; when controlling for the size of other brain structures, cerebellum size correlates markedly more strongly with foraging skill than it does with social group size and more strongly than neocortex size does with foraging skill, whereas for neocortex size the reverse pattern is observed. This is confirmed by analyses of each structure with the other included as a predictor; neocortex size then correlates significantly with social group size ($t_{6,36} = 3.92$, $p = 0.0005$) but not extractive foraging ($t_{6,36} = 1.01$, $p = 0.32$), whereas cerebellum size correlates significantly with extractive foraging ($t_{6,36} = 3.59$, $p = 0.001$) but not social group size ($t_{6,36} = 1.33$, $p = 0.19$). Although these results, together with those showing cerebellum-specific expansion in apes, certainly imply a degree of functional dissociation and independent evolution of the two structures, it is important to emphasize that each structure does correlate with both behavioural

variables when not controlling for the other (in line with the evidence of coordinated cortico-cerebellar evolution). Thus, behavioural specializations seem to be based on a combination of both independent and coordinated evolution of individual brain structures.

Primate tool use frequently occurs in the context of extractive foraging and involves similarly complex, organized sequences of behaviours [113]. Fewer species are recorded as using tools than using extractive foraging [7]. Nevertheless, broadly similar results are obtained for tool use. Controlling for body size, and residual brain volume, cerebellum size correlates with tool use ($t_{5,36} = 2.04$, $p = 0.050$) but not social group size ($t_{6,36} = 1.47$, $p = 0.15$), while neocortex size correlates with social group size ($t_{6,36} = 3.98$, $p = 0.0003$) but not tool use ($t_{6,36} = 0.71$, $p = 0.48$).

9. CO-EVOLUTION OF SOCIAL AND TECHNICAL INTELLIGENCE

The debate about whether it was selection on social or technical intelligence that drove the evolution of brain size and cognitive capacities has increasingly appeared to be resolved in favour of the former [8,9]. Based on the evidence presented above, and in common with some other recent authors [33,108,112–114], I suggest not only that selection pressures on both social and technical skills were important, but also that they interacted with one another during human evolution. The theoretical argument is elaborated by Barrett *et al.* [33], who persuasively argue that the social and physical environment form mutually reinforcing feedback loops.

Specialization for technical intelligence seems particularly relevant to aspects of great ape behaviour. Great apes do not live in particularly large groups, but they are adept at extractive foraging and tool use, and at learning these skills by observation of others [104,105,113]. The capacities to perform such behaviours, and to learn them by observing others, may be intrinsically linked. Byrne [112] argues that both depend on ‘behaviour parsing’: the capacity to segment and mentally organize a sequence of acts into its subroutines based on the statistical regularities among the observed acts. This capacity is likely to have its origin in foraging skills: the relative lack of physiological adaptations for digesting high-fibre plant material in apes compared to Old World monkeys would have put a premium on extraction of more nutritious resources from hard or tough shells, spiny plants, termite mounds and other challenging defences. Once, however, the capacity to parse action sequences was established, it could have been secondarily adapted for use in the social domain, forming a basis for the prediction of conspecifics’ behaviour [108–112].

10. EMBODIED SIMULATION AND SOCIAL UNDERSTANDING

A sensory–motor origin of socio-cognitive capacities, and a linkage between the ability to execute complex behavioural sequences and to perceive and decode them when observing others, both fit with data indicating that the neural systems activated during a

particular behaviour are also activated when observing the same behaviour performed by another individual [117]. It may therefore be that simulating the neural states underlying behaviours contributes to understanding them during observation. For example, the recognition of emotional expressions is disrupted by transcranial magnetic stimulation of somatosensory cortex, implying that activation of the system for producing expressions contributes to decoding them [118]. Computational work also supports the idea that simulation may provide a direct link between sensory–motor control and social understanding [119], and there are close computational parallels between motor control and control of social interactions [120].

Although most work on embodied social simulation has focused on the activity of ‘mirror neurons’ localised to a few cortical regions, such mirror-like properties are likely to be a function of the way that neurons are embedded in more distributed neural networks involved in sensory–motor processing [121–124], and experimental evidence now implicates the cerebellum [85–87,90,125,126]. The ‘mirror neuron system’ may thus not be a functionally specialized neural circuit restricted to a few cortical areas, nor an adaptation evolved specifically for action understanding, and as such may not merit the term ‘system’ [121]. Instead, mirroring may be a rather general adaptive property of neural systems with the right architecture for forming associations between one’s own and others’ actions, and may be phylogenetically widespread [127].

Damasio and Meyer [123] outline in broad form a model of mirror neurons based on ‘retro-activation’, the key to which is a neural architecture in which anterior association areas send signals back to visual cortex (and even to the visual thalamus). The comparatively large size and great complexity of primate visual and visuo–motor systems, including numerous reciprocal connections between anterior and posterior visual areas, and between these areas and association areas in frontal and temporal cortices [68,128], may therefore have implications for primate social cognition without necessarily having evolved primarily as an adaptation for it. However, an interesting question is then whether, once a sensory–motor system has mirroring potential, this potential is exploited by further evolutionary adaptive strengthening of critical connections in more social species, or perhaps inhibited in species or domains of behaviour where mirroring would be disadvantageous (for example, mirroring of subordinate expressions in dominance interactions).

11. CONCLUSIONS

The search for a single ideal comparative brain measure that captures the neural basis of cognitive evolution is likely to be more obfuscatory than illuminating, because different selection pressures have acted on different neural systems at different times. Whilst there are general patterns, such as the tendency of neocortex and cerebellum to evolve together, there are also significant deviations from such trends, such as visual pathway expansion in primates, and cerebellar expansion in apes. Gross brain size and composite brain indices or ratios therefore conflate different

neural adaptations and mask important evolutionary patterns. To understand the neural bases of cognitive evolution, appropriate statistical, phylogenetic analyses that tease apart the variation associated with different neural systems and due to different selection pressures will therefore be more useful than composite indices.

Any account of human neuro-cognitive evolution needs to explain why there are so many neurons in the cerebellum. The answer suggested here, based on converging comparative and experimental evidence, is that the cerebellum and cortico-cerebellar networks are key components of systems enabling the control, organization and comprehension of complex sequences involved in both technical and social intelligence, and, ultimately, language. These proposals agree with Sterelny's [114] scenario for language evolution which suggests that the control of motor sequences involved in ape foraging skills provided a cognitive platform for gestural communication and thence ultimately syntax and language, and with Fitch's [101] proposal that motor control and hierarchical action planning systems were secondarily adapted for syntax.

The evidence presented here suggests that sensory-motor and cognitive evolution are not dissociable. In common with Barrett [33], I argue that there is no need to postulate a distinct set of 'cognitive' processes to fill the supposed gap between sensory reception and motor output. Even 'offline' and seemingly abstract cognitive processes, such as number representation and metaphor, appear to be 'body based' [31,129], and many allegedly abstract, centralized cognitive processes recruit distributed sensory-motor systems that evolved to control bodily movement [31]. By extension, cognitive evolution is to be understood as the elaboration of embodied control systems, rather than of a disembodied reasoning device [28,30]. As a corollary, there is no 'intelligent', 'executive' or indeed 'Fodorian' [130] bit of the brain that holds the key to cognitive evolution. Instead, the evolution of large brains was associated with the elaboration of sensory-motor mechanisms for the adaptive control of bodies in their environments.

I thank Celia Heyes, Russell Gray, Kim Sterelny, Eva Jablonka, Alison Gopnik, Arthur Robson, Matthew Rushworth and Nick Shea for many useful discussions, Simon Reader for access to extractive foraging and tool use data from ref. 7, and Andy Whiten, Dick Byrne and two anonymous referees for additional comments on the manuscript.

REFERENCES

- Harvey, P. H. & Krebs, J. R. 1990 Comparing brains. *Science* **249**, 140–146. (doi:10.1126/science.2196673)
- Shettleworth, S. J. 1998 *Cognition, evolution, and behavior*. New York, NY: Oxford University Press.
- Striedter, G. F. 2005 *Principles of brain evolution*. Sunderland, MA: Sinauer.
- Dunbar, R. I. M. & Shultz, S. 2007 Evolution in the social brain. *Science* **317**, 1344–1347. (doi:10.1126/science.1145463)
- Sol, D. 2009 Revisiting the cognitive buffer hypothesis for the evolution of large brains. *Biol. Lett.* **5**, 130–133. (doi:10.1098/rsbl.2008.0621)
- Lefebvre, L. & Sol, D. 2008 Brains, lifestyles and cognition: are there general trends? *Brain Behav. Evol.* **72**, 135–144. (doi:10.1159/000151473)
- Reader, S. M., Hager, Y. & Laland, K. N. 2011 The evolution of primate general and cultural intelligence. *Phil. Trans. R. Soc. B* **366**, 1017–1027. (doi:10.1098/rstb.2010.0342)
- Dunbar, R. I. M. 1998 The social brain hypothesis. *Evol. Anthropol.* **6**, 178–190. (doi:10.1002/(SICI)1520-6505(1998)6:5<178::AID-EVAN5>3.0.CO;2-8)
- Dunbar, R. I. M. & Shultz, S. 2007 Understanding primate brain evolution. *Phil. Trans. R. Soc. B* **362**, 649–658. (doi:10.1098/rstb.2006.2001)
- Keverne, E. B., Martel, F. L. & Nevison, C. M. 1996 Primate brain evolution: genetic and functional considerations. *Proc. R. Soc. Lond. B* **263**, 689–696. (doi:10.1098/rspb.1996.0103)
- Reader, S. M. & Laland, K. N. 2002 Social intelligence, innovation, and enhanced brain size in primates. *Proc. Natl Acad. Sci. USA* **99**, 4436–4441. (doi:10.1073/pnas.062041299)
- Lefebvre, L., Reader, S. M. & Sol, D. 2004 Brains, innovations and evolution in birds and primates. *Brain Behav. Evol.* **63**, 233–246. (doi:10.1159/000076784)
- Deaner, R. O., Isler, K., Burkhardt, J. & van Schaik, C. P. 2007 Overall brain size, and not encephalization quotient, best predicts cognitive ability across non-human primates. *Brain Behav. Evol.* **70**, 115–124. (doi:10.1159/000102973)
- Healy, S. D. & Rowe, C. 2007 A critique of comparative studies of brain size. *Proc. R. Soc. B* **274**, 453–464. (doi:10.1098/rspb.2006.3748)
- Krebs, J. R. 1990 Food-storing birds: adaptive specialization in brain and behaviour? *Phil. Trans. R. Soc. Lond. B* **329**, 153–160. (doi:10.1098/rstb.1990.0160)
- Barton, R. A. & Dean, P. 1993 Comparative evidence indicating neural specialization for predatory behaviour in mammals. *Proc. R. Soc. Lond. B* **254**, 63–68. (doi:10.1098/rspb.1993.0127)
- Devoogd, T. J., Krebs, J. R., Healy, S. D. & Purvis, A. 1993 Relations between song repertoire size and the volume of brain nuclei related to song: comparative evolutionary analyses amongst oscine birds. *Proc. R. Soc. Lond. B* **254**, 75–82. (doi:10.1098/rspb.1993.0129)
- Barton, R. A., Purvis, A. & Harvey, P. H. 1995 Evolutionary radiation of visual and olfactory brain systems in primates, bats and insectivores. *Phil. Trans. R. Soc. Lond. B* **348**, 381–392. (doi:10.1098/rstb.1995.0076)
- Barton, R. A. 1998 Visual specialisation and brain evolution in primates. *Proc. R. Soc. Lond. B* **265**, 1933–1937. (doi:10.1098/rspb.1998.0523)
- Barton, R. A. 2004 Binocularity and brain evolution in primates. *Proc. Natl Acad. Sci. USA* **101**, 10 113–10 115. (doi:10.1073/pnas.0401955101)
- Iwaniuk, A. N. & Hurd, P. L. 2005 The evolution of cerebrotypes in birds. *Brain Behav. Evol.* **65**, 215–230. (doi:10.1159/000084313)
- Shumway, C. A. 2010 The evolution of complex brains and behaviors in African cichlid fishes. *Curr. Zool.* **56**, 144–156.
- Barton, R. A. & Harvey, P. H. 2000 Mosaic evolution of brain structure in mammals. *Nature* **405**, 1055–1058. (doi:10.1038/35016580)
- De Winter, W. & Oxnard, C. E. 2001 Evolutionary radiations and convergences in the structural organization of mammalian brains. *Nature* **409**, 710–714. (doi:10.1038/35055547)
- Herculano-Houzel, S. 2009 The human brain in numbers: a linearly scaled-up primate brain. *Front Neurosci.* **3**, 31.
- Herculano-Houzel, S. 2010 Coordinated scaling of cortical and cerebellar numbers of neurons. *Front. Neuroanat.* **4**, 12. (doi:10.3389/fnana.2010.00012)

- 27 Fodor, J. 1983 *The modularity of mind*. Cambridge, MA: MIT Press.
- 28 Damasio, A. 1994 *Descartes' error: emotion, reason, and the human brain*. New York, NY: Putnam.
- 29 Chiel, H. J. & Beer, R. D. 1997 The brain has a body: adaptive behavior emerges from interactions of nervous system, body and environment. *Trends Neurosci.* **20**, 553–557. (doi:10.1016/S0166-2236(97)01149-1)
- 30 Clark, A. 1997 *Being there: putting brain, body, and world together again*. Cambridge, MA: MIT Press.
- 31 Wilson, M. 2002 Six views of embodied cognition. *Psychol. Bull. Rev.* **9**, 625–636. (doi:10.3758/BF03196322)
- 32 Anderson, M. L. 2010 Neural reuse: a fundamental organizational principle of the brain. *Behav. Brain Sci.* **33**, 245–313. (doi:10.1017/S0140525X10000853)
- 33 Barrett, L., Henzi, S. P. & Lusseau, D. 2012 Taking sociality seriously: the structure of multi-dimensional social networks as a source of information for individuals. *Phil. Trans. R. Soc. B* **367**, 2108–2118. (doi:10.1098/rstb.2012.0113)
- 34 Gallese, V. & Lakoff, G. 2005 The brain's concepts: the role of the sensory-motor system in conceptual knowledge. *Cogn. Neuropsychol.* **22**, 455–479. (doi:10.1080/02643290442000310)
- 35 Nunn, C. L. 2011 *The comparative approach in evolutionary anthropology and biology*. Chicago, IL: Chicago University Press.
- 36 Pagel, M. 1999 The maximum likelihood approach to reconstructing ancestral character states of discrete characters on phylogenies. *Syst. Biol.* **48**, 612–622. (doi:10.1080/106351599260184)
- 37 Rohlf, F. J. 2001 Comparative methods for the analysis of continuous variables: geometric interpretations. *Evolution* **55**, 2143–2160.
- 38 Freckleton, R. P., Harvey, P. H. & Pagel, M. 2002 Phylogenetic analysis and comparative data: a test and review of evidence. *Am. Nat.* **160**, 712–726. (doi:10.1086/343873)
- 39 Lui, J. H., Hansen, D. V. & Kriegstein, A. R. 2011 Development and evolution of the human neocortex. *Cell* **146**, 18–36. (doi:10.1016/j.cell.2011.06.030)
- 40 Rakic, P. 2009 Evolution of the neocortex: a perspective from developmental biology. *Nat. Rev. Neurosci.* **12**, 724–735. (doi:10.1038/nrn2719)
- 41 Carlson, K. J., Stout, D., Jashashvili, T., de Ruiter, D. J., Tafforeau, P., Carlson, K. & Berger, L. R. 2011 The endocast of MH1, *Australopithecus sediba*. *Science* **333**, 1402–1407. (doi:10.1126/science.1203922)
- 42 Stephan, H., Frahm, H. D. & G, Baron 1981 New and revised data on volumes of brain structures in insectivores and primates. *Folia Primatol.* **35**, 1–29. (doi:10.1159/000155963)
- 43 Stephan, H., Baron, G. & Frahm, H. D. 1991 *Comparative brain research in mammals*, vol. 1. *Insectivores*. New York, NY: Springer.
- 44 Bush, E. C. & Allman, J. M. 2004 The scaling of frontal cortex size in primates and carnivores. *Proc. Natl Acad. Sci. USA* **101**, 3962–3966. (doi:10.1073/pnas.0305760101)
- 45 Mangold-Wirz, K. 1966 Cerebralisierung und Ontogenese bei Eutheriern. *Acta Anat.* **63**, 449. (doi:10.1159/000142809)
- 46 Zhang, K. & Sejnowski, T. J. 2000 A universal scaling law between gray matter and white matter of cerebral cortex. *Proc. Natl Acad. Sci. USA* **97**, 5621–5626. (doi:10.1073/pnas.090504197)
- 47 Bush, E. C. & Allman, J. M. 2003 The scaling of white matter to grey matter in cerebellum and neocortex. *Brain Behav. Evol.* **61**, 1–5. (doi:10.1159/000068880)
- 48 Barton, R. A. 2006 Primate brain evolution: integrating comparative, neurophysiological and ethological data. *Evol. Anthropol.* **15**, 224–236. (doi:10.1002/evan.20105)
- 49 Wang, S.-H., Shultz, J. R., Burish, M. J., Harrison, K. H., Hof, P. R., Towns, L. C., Wagers, M. W. & Wyatt, K. D. 2008 Functional trade-offs in white matter axonal scaling. *J. Neurosci.* **28**, 4047–4056. (doi:10.1523/JNEUROSCI.5559-05.2008)
- 50 Clark, D. A., Mitra, P. P. & Wang, S. S. H. 2001 Scalable architecture in mammalian brains. *Nature* **411**, 189–193. (doi:10.1038/35075564)
- 51 Brodmann, K. 1912 Neue Ergebnisse über die vergleichende histologische Lokalisation der Grosshirnrinde mit besonderer Berücksichtigung des Stirnhirns. *Anat. Anzeiger* **41**, 157–216.
- 52 Blinkov, S. M. & Glezer, I. I. 1968 *Das Zentralnervensystem in Zahlen und Tabellen*. Jena, Germany: Fischer.
- 53 Rilling, J. K. 2006 Human and non-human primate brains: are they allometrically scaled versions of the same design? *Evol. Anthropol.* **15**, 65–77. (doi:10.1002/evan.20095)
- 54 Semendeferi, K., Lu, A., Schenker, N. & Damasio, H. 2002 Humans and great apes share a large frontal cortex. *Nat. Neurosci.* **5**, 272–276. (doi:10.1038/nn814)
- 55 Schoenemann, P. T., Sheehan, M. J. & Glotzer, L. D. 2005 Prefrontal white matter volume is disproportionately larger in humans than in other primates. *Nat. Neurosci.* **8**, 242–252. (doi:10.1038/nn1394)
- 56 Sherwood, C. C., Holloway, R. L., Semendeferi, K. & Hof, P. R. 2005 Is prefrontal white matter enlargement a human evolutionary specialization? *Nat. Neurosci.* **8**, 537–538. (doi:10.1038/nn0505-537)
- 57 Smaers, J. B., Steele, J., Case, C. R., Cowper, A., Amunts, K. & Zilles, K. 2011 Primate prefrontal cortex evolution: human brains are the extreme of a lateralized ape trend. *Brain Behav. Evol.* **77**, 67–78. (doi:10.1159/000323671)
- 58 Semendeferi, K., Teffer, K., Buxhoeveden, D. P., Park, M. S., Bludau, S., Amunts, K., Travis, K. & Buckwalter, J. 2011 Spatial organization of neurons in the frontal pole sets humans apart from great apes. *Cereb. Cortex* **21**, 1485–1497. (doi:10.1093/cercor/bhq191)
- 59 Dunbar, R. I. M. & Shultz, S. 2007 Understanding primate brain evolution. *Phil. Trans. R. Soc. B* **362**, 649–658. (doi:10.1098/rstb.2006.2001)
- 60 Semendeferi, K., Armstrong, E., Schleicher, A., Zilles, K. & Van Hoesen, G. W. 2001 Prefrontal cortex in humans and apes: a comparative study of area 10. *Am. J. Phys. Anthropol.* **114**, 224–241. (doi:10.1002/1096-8644(200103)114:3<224::AID-AJPA1022>3.0.CO;2-I)
- 61 Balsters, J. H., Cussans, E., Diedrichsen, J., Phillips, K. A., Preuss, T. M., Rilling, J. K. & Ramnani, N. 2009 Evolution of the cerebellar cortex: the selective expansion of prefrontal-projecting cerebellar lobules. *NeuroImage* **49**, 2045–2052. (doi:10.1016/j.neuroimage.2009.10.045)
- 62 Smaers, J. B., Steele, J. & Zilles, K. 2011 Modeling the evolution of cortico-cerebellar systems in primates. *Ann. NY Acad. Sci.* **1225**, 176–190. (doi:10.1111/j.1749-6632.2011.06003.x)
- 63 Barton, R. A. & Venditti, C. Submitted. Human frontal lobes are not disproportionately expanded. *Nat. Neurosci.*
- 64 Rilling, J. K. & Seligman, R. A. 2002 A quantitative morphometric comparative analysis of the primate temporal lobe. *J. Hum. Evol.* **42**, 505–533. (doi:10.1006/jhev.2001.0537)
- 65 Barger, N., Stefanacci, L. & Semendeferi, K. 2007 A comparative volumetric analysis of the amygdaloid

- complex and basolateral division in the human and ape brain. *Am. J. Phys. Anthropol.* **134**, 392–403. (doi:10.1002/ajpa.20684)
- 66 Barton, R. A. 2000 Ecological and social factors in primate brain evolution. In *On the move: how and why animals travel in groups* (eds S. Boinski & P. Garber), pp. 204–237. Chicago, IL: Chicago University Press.
- 67 Whiting, B. & Barton, R. A. 2003 The evolution of the cortico-cerebellar complex in primates: anatomical connections predict patterns of correlated evolution. *J. Hum. Evol.* **44**, 3–10. (doi:10.1016/S0047-2484(02)00162-8)
- 68 van Essen, D. C., Anderson, C. H. & Felleman, D. J. 1992 Information processing in the primate visual system: an integrated systems perspective. *Science* **255**, 419–423. (doi:10.1126/science.1734518)
- 69 Rilling, J. K. & Insel, T. R. 1998 Evolution of the cerebellum in primates: differences in relative volume among monkeys, apes and humans. *Brain Behav. Evol.* **52**, 308–314. (doi:10.1159/00006575)
- 70 MacLeod, C. E., Zilles, K., Schleicher, A., Rilling, J. K. & Gibson, K. R. 2003 Expansion of the neocerebellum in Hominoidea. *J. Hum. Evol.* **44**, 401–429. (doi:10.1016/S0047-2484(03)00028-9)
- 71 Barton, R. A. & Venditti, C. In revision. Explosive evolution of the cerebellum in humans and other great apes. *Curr. Biol.*
- 72 Glickstein, M. & Doron, K. 2008 Cerebellum: connections and functions. *Cerebellum* **7**, 589–594. (doi:10.1007/s12311-008-0074-4)
- 73 Habas, C. 2010 Functional imaging of the deep cerebellar nuclei: a review. *Cerebellum* **9**, 22–28. (doi:10.1007/s12311-009-0119-3)
- 74 Thier, P. & Ilg, U. J. 2005 The neural basis of smooth-pursuit eye movements. *Curr. Opin. Neurobiol.* **15**, 645–652. (doi:10.1016/j.conb.2005.10.013)
- 75 Colibazzi, T. *et al.* 2010 Neural systems subserving valence and arousal during the experience of induced emotions. *Emotion* **10**, 377–389. (doi:10.1037/a0018484)
- 76 Tavano, A. & Borgatti, R. 2010 Evidence for a link among cognition, language and emotion in cerebellar malformations. *Cortex* **46**, 907–918. (doi:10.1016/j.cortex.2009.07.017)
- 77 Bellebaum, C. & Daum, I. 2011 Mechanisms of cerebellar involvement in associative learning. *Cortex* **47**, 128–136. (doi:10.1016/j.cortex.2009.07.016)
- 78 Leiner, H. C. 2010 Solving the mystery of the human cerebellum. *Neuropsychol. Rev.* **20**, 229–235. (doi:10.1007/s11065-010-9140-z)
- 79 Schmahmann, J. D. & Sherman, J. C. 1998 The cerebellar cognitive affective syndrome. *Brain* **121**, 561–579. (doi:10.1093/brain/121.4.561)
- 80 Steinlin, M. 2008 Cerebellar disorders in childhood: cognitive problems. *Cerebellum* **7**, 607–610. (doi:10.1007/s12311-008-0083-3)
- 81 Leggio, M. G., Chiricozzi, F. R., Clausi, S., Tedesco, A. M. & Molinari, M. 2011 The neuropsychological profile of cerebellar damage: the sequencing hypothesis. *Cortex* **47**, 137–144. (doi:10.1016/j.cortex.2009.08.011)
- 82 Rochefort, C., Arabo, A., André, M., Poucet, B., Save, E. & Rondi-Reig, L. 2011 Cerebellum shapes hippocampal spatial code. *Science* **334**, 385–389. (doi:10.1126/science.1207403)
- 83 Forster, S. E. & Brown, J. W. 2011 Medial prefrontal cortex predicts and evaluates the timing of action outcomes. *NeuroImage* **55**, 253–265. (doi:10.1016/j.neuroimage.2010.11.035)
- 84 Ramnani, N. & Miall, C. R. 2004 A system in the human brain for predicting the actions of others. *Nat. Neurosci.* **7**, 85. (doi:10.1038/nn1168)
- 85 Gazzola, V. & Keysers, C. 2009 The observation and execution of actions share motor and somatosensory voxels in all tested subjects: single-subject analyses of unsmoothed fMRI data. *Cereb. Cortex* **19**, 1239–1255. (doi:10.1093/cercor/bhn181)
- 86 Schulte-Ruther, M., Markowitsch, H. J., Fink, G. R. & Piefke, M. 2007 Mirror neuron and theory of mind mechanisms involved in face-to-face interactions: a functional magnetic resonance imaging approach to empathy. *J. Cogn. Neurosci.* **19**, 1354–1372. (doi:10.1162/jocn.2007.19.8.1354)
- 87 Singer, T., Seymour, B., O’Doherty, J., Kaube, H., Dolan, R. J. & Frith, C. D. 2004 Empathy for pain involves affective but not sensory components of pain. *Science* **303**, 1157–1162. (doi:10.1126/science.1093535)
- 88 Jackson, P. L., Meltzoff, A. N. & Decety, J. 2005 Neural circuits involved in imitation and perspective-taking. *NeuroImage* **31**, 429–439. (doi:10.1016/j.neuroimage.2005.11.026)
- 89 Ito, M. 2008 Control of mental activities by internal models in the cerebellum. *Nat. Rev. Neurosci.* **9**, 304–313. (doi:10.1038/nrn2332)
- 90 Strick, P. L., Dum, R. P. & Fiez, J. A. 2009 Cerebellum and nonmotor function. *Annu. Rev. Neurosci.* **32**, 413–434. (doi:10.1146/annurev.neuro.31.060407.125606)
- 91 Hogan, M. J., Staff, R. T., Bunting, B. P., Murray, A. D., Ahearn, T. S., Deary, I. J. & Whalley, L. J. 2011 Cerebellar brain volume accounts for variance in cognitive performance in older adults. *Cortex* **47**, 441–450. (doi:10.1016/j.cortex.2010.01.001)
- 92 Shukla, D. K., Keehn, B., Lincoln, A. J. & Muller, R. A. 2010 White matter compromise of callosal and subcortical fiber tracts in children with autism spectrum disorder: a diffusion tensor imaging study. *J. Am. Acad. Child Adol. Psych.* **49**, 1269–1278.
- 93 Glickstein, M., Sultan, F. & Voogd, J. 2008 Functional localization in the cerebellum. *Cortex* **47**, 59–80. (doi:10.1016/j.cortex.2009.09.001)
- 94 Beaton, A. & Mariën, P. 2010 Language, cognition and the cerebellum: grappling with an enigma. *Cortex* **46**, 811–820. (doi:10.1016/j.cortex.2010.02.005)
- 95 Ramnani, N. 2006 The primate cortico-cerebellar system: anatomy and function. *Nat. Rev. Neurosci.* **7**, 511. (doi:10.1038/nrn1953)
- 96 Ito, M. 1993 Movement and thought: identical control mechanisms by the cerebellum. *Trends Neurosci.* **16**, 448–450. (doi:10.1016/0166-2236(93)90073-U)
- 97 Hickok, G. 2012 Computational neuroanatomy of speech production. *Nat. Rev. Neurosci.* **13**, 135–145. (doi:10.1038/nrg3118)
- 98 Behrens, T. E., Hunt, L. T. & Rushworth, M. F. 2009 The computation of social behavior. *Science* **324**, 1160–1164. (doi:10.1126/science.1169694)
- 99 Krossa, E., Bermanna, M. G., Mischel, W., Smith, E. E. & Wager, T. D. 2011 Social rejection shares somatosensory representations with physical pain. *Proc. Natl Acad. Sci. USA* **108**, 6270–6275. (doi:10.1073/pnas.1102693108)
- 100 Shackman, A. J., Salomons, T. V., Slagter, H. A., Fox, A. S., Winter, J. J. & Davidson, R. J. 2011 The integration of negative affect, pain and cognitive control in the cingulate cortex. *Nat. Rev. Neurosci.* **12**, 154–167. (doi:10.1038/nrn2994)
- 101 Fitch, W. T. 2011 The evolution of syntax: an exaptationist perspective. *Front. Evol. Neurosci.* **3**, 9. (doi:10.3389/fnevo.2011.00009)
- 102 Imamizu, H., Miyauchi, S., Tamada, T., Sasaki, Y., Takino, R., Putz, B., Yoshioko, T. & Kawato, M. 2000 Human cerebellar activity reflecting an acquired internal model of a new tool. *Nature* **403**, 192–195. (doi:10.1038/35003194)

- 103 Mar, R. A. 2011 The neural bases of social cognition and story comprehension. *Annu. Rev. Psychol.* **62**, 103–134. (doi:10.1146/annurev-psych-120709-145406)
- 104 Whiten, A., Goodall, J., McGrew, W. C., Nishida, T., Reynolds, V., Sugiyama, Y., Tutin, C. E., Wrangham, R. W. & Boesch, C. 1999 Cultures in chimpanzees. *Nature* **399**, 682–685. (doi:10.1038/21415)
- 105 Whiten, A. & Suddendorf, T. 2007 Great ape cognition and the evolutionary roots of human imagination. *Proc. Br. Acad.* **147**, 31–59. (doi:10.5871/bacad/9780197264195.001.0001)
- 106 Endress, A. D., Carden, S., Versace, E. & Hauser, M. D. 2010 The apes' edge: positional learning in chimpanzees and humans. *Anim. Cogn.* **13**, 483–495. (doi:10.1007/s10071-009-0299-8)
- 107 Byrne, R. W. 1999 Object manipulation and skill organization in the complex food preparation of mountain gorillas. In *The mentality of gorillas and orangutans* (eds S. T. Parker, R. W. Mitchell & H. L. Miles), pp. 147–159. Cambridge, UK: Cambridge University Press. (doi:10.1017/CBO9780511542305.007)
- 108 Byrne, R. W. & Bates, L. A. 2010 Primate social cognition: uniquely primate, uniquely social, or just unique? *Neuron* **65**, 815–830. (doi:10.1016/j.neuron.2010.03.010)
- 109 Byrne, R. W., Hobaiter, C. & Klailova, M. 2011 Local traditions in gorilla manual skill: evidence for observational learning of behavioral organization. *Anim. Cogn.* **14**, 683–693. (doi:10.1007/s10071-011-0403-8)
- 110 Byrne, R. W. 2007 Culture in great apes: using intricate complexity in feeding skills to trace the evolutionary origin of human technical prowess. *Phil. Trans. R. Soc. B* **362**, 577–585. (doi:10.1098/rstb.2006.1996)
- 111 Stout, D., Toth, N., Schick, K. & Chaminade, T. 2008 Neural correlates of Early Stone Age toolmaking: technology, language and cognition in human evolution. *Phil. Trans. R. Soc. B* **363**, 1939–1949. (doi:10.1098/rstb.2008.0001)
- 112 Byrne, R. W. 2006 Parsing behaviour. A mundane origin for an extraordinary ability? In *The roots of human sociality* (eds S. Levinson & N. Enfield), pp. 478–505. Oxford, UK: Berg.
- 113 Stout, D., Passingham, R., Frith, C., Apel, J. & Chaminade, T. 2011 Technology, expertise and social cognition in human evolution. *Eur. J. Neurosci.* **33**, 1328–1338. (doi:10.1111/j.1460-9568.2011.07619.x)
- 114 Sterelny, K. 2012 Language, gesture, skill: the co-evolutionary foundations of language. *Phil. Trans. R. Soc. B* **367**, 2141–2151. (doi:10.1098/rstb.2012.0116)
- 115 Londei, A., D'Ausilio, A., Basso, D., Sestieri, C., Del Gratta, C., Romani, G. L. & Belardinelli, M. O. 2010 Sensory-motor brain network connectivity for speech comprehension. *Hum. Brain Mapp.* **4**, 567–580.
- 116 Gibson, K. R. 1986 Cognition, brain size and the extraction of embedded food resources. In *Primate ontogeny, cognition and social behaviour*. (eds J. G. Else & P. C. Lee), pp. 93–103. Cambridge, UK: Cambridge University Press.
- 117 Gallese, V. 2009 Motor abstraction: a neuroscientific account of how action goals and intentions are mapped and understood. *Psychol. Res.* **73**, 486–498. (doi:10.1007/s00426-009-0232-4)
- 118 Pitcher, D., Garrido, L., Walsh, V. & Duchaine, B. C. 2008 Transcranial magnetic stimulation disrupts the perception and embodiment of facial expressions. *J. Neurosci.* **28**, 8929–8933. (doi:10.1523/JNEUROSCI.1450-08.2008)
- 119 Oztop, E., Wolpert, D. & Kawato, M. 2005 Mental state inference using visual control parameters. *Cog. Brain Res.* **22**, 129–151. (doi:10.1016/j.cogbrainres.2004.08.004)
- 120 Wolpert, D. M., Doya, K. & Kawato, M. 2003 A unifying computational framework for motor control and social interaction. *Phil. Trans. R. Soc. Lond. B* **358**, 593–602. (doi:10.1098/rstb.2002.1238)
- 121 Heyes, C. M. 2010 Where do mirror neurons come from? *Neurosci. Biobehav. Rev.* **34**, 575–583. (doi:10.1016/j.neubiorev.2009.11.007)
- 122 Keysers, C. & Perrett, D. I. 2004 Demystifying social cognition: a Hebbian perspective. *Trends Cogn. Sci.* **8**, 501–507. (doi:10.1016/j.tics.2004.09.005)
- 123 Damasio, A. & Meyer, K. 2008 Behind the looking-glass. *Nature* **454**, 167–168. (doi:10.1038/454167a)
- 124 Gallese, V. 2008 Mirror neurons and the social nature of language: the neural exploitation hypothesis. *Soc. Neurosci.* **3**, 317–333. (doi:10.1080/17470910701563608)
- 125 Peeters, R., Simone, L., Nelissen, K., Fabbri-Destro, M., Vanduffel, W., Rizzolatti, G. & Orban, G. A. 2009 The representation of tool use in humans and monkeys: common and uniquely human features. *J. Neurosci.* **29**, 11 523–11 539.
- 126 Kana, R. K., Wadsworth, H. M. & Travers, B. G. 2011 A systems level analysis of the mirror neuron hypothesis and imitation impairments in autism spectrum disorders. *Neurosci. Biobehav. Rev.* **35**, 894–902. (doi:10.1016/j.neubiorev.2010.10.007)
- 127 Bonini, L. & Ferrari, P. F. 2011 Evolution of mirror systems: a simple mechanism for complex cognitive functions. *Ann. NY Acad. Sci.* **1225**, 166–175. (doi:10.1111/j.1749-6632.2011.06002.x)
- 128 Young, M. P. 2000 The architecture of visual cortex and inferential processes in vision. *Spat. Vis.* **13**, 137–146. (doi:10.1163/156856800741162)
- 129 Santana, E. & de Vega, M. 2011 Metaphors are embodied, and so are their literal counterparts. *Front. Psychol.* **2**, 90. (doi:10.3389/fpsyg.2011.00090)
- 130 Bolhuis, J. J., Brown, G. R., Richardson, R. C. & Laland, K. N. 2011 Darwin in mind: new opportunities for evolutionary psychology. *PLoS Biol.* **9**, e1001109. (doi:10.1371/journal.pbio.1001109)